

Anti-Human β -NGF Antibody
Catalog # ABG10524**Specification**

Anti-Human β -NGF Antibody - Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Mouse
Clonality	Monoclonal

Anti-Human β -NGF Antibody - Additional Information**Preparation**

Produced in mice using highly pure recombinant human β -NGF as the immunizing antigen. This IgG2_{bK} antibody was purified from cell culture by Protein G affinity chromatography.

WesternBlot

To detect Human β -NGF by Western Blot analysis this antibody can be used at a concentration of 4.0-8.0 μ g/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human β -NGF is 4.0-8.0 ng/lane, under reducing or non-reducing conditions.

Sandwich

In a sandwich ELISA (assuming 100 μ l/well), a concentration of 2.0-4.0 μ g/ml of this antibody will detect at least 200 pg/ml of recombinant Human β -NGF when used with BioGems's biotinylated antigen affinity purified anti-Human β -NGF (60-247BT) as the detection antibody at a concentration of approximately 0.5-1.0 μ g/ml.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of human pancreas adenocarcinoma. The recommended concentration is 2.5 μ g/mL with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 9.0 buffer is recommended. See photo(s). Optimal concentrations and conditions may vary.

 Tissue samples were provided by the Cooperative Human Tissue Network, which is funded by the National Cancer Institute. Information and photo are courtesy of Flagship Biosciences, LLC - Histology Services.

Neutralization

To yield one-half maximal inhibition [ND₅₀] of the biological activity of Human β -NGF (10.0 ng/ml), a concentration of 2.7 - 4.0 μ g/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human β -NGF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human β -NGF Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human β -NGF Antibody - Images