

Anti-Human TECK Antibody

Catalog # ABG10535

Specification

Anti-Human TECK Antibody - Product Information

Application WB, IHC, E
Reactivity Human
Host Mouse
Clonality Monoclonal

Anti-Human TECK Antibody - Additional Information

Preparation

Produced in BALB/c x ICR F₁ mice using highly pure (>98%) recombinant human TECK as the immunizing antigen. This IgG1_K antibody was purified from ascites fluid by Protein A affinity chromatography.

WesternBlot

To detect hTECK by Western Blot analysis this antibody can be used at a concentration of 0.25- $0.50 \mu g/ml$. Used in conjunction with compatible secondary reagents the detection limit for recombinant hTECK is 0.50- $1.0 \mu g/lane$, under non-reducing conditions.

Sandwich

In a sandwich ELISA (assuming 100μ l/well), a concentration of 1.0- $2.0~\mu$ g/ml of this antibody will detect at least 300 pg/ml of recombinant human TECK when used with BioGems' biotinylated antigen affinity purified anti-human TECK (60-305BT) as the detection antibody at a concentration of approximately 0.5- $1.0~\mu$ g/ml.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of human normal small bowel. The recommended concentration is 2.5 μ g/mL with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen and a Proteinase K antigen retrieval. Optimal concentrations and conditions may vary.

 >lnformation and photo are courtesy of Flagship Biosciences, LLC - Histology Services.

Formulation

A sterile filtered antibody solution was lyophilized from PBS.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human TECK Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



Anti-Human TECK Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Human TECK Antibody - Images