

**Goat Anti-HOXA10 Antibody (internal region), Biotinylated**  
**Catalog # AF4299a****Specification**

---

**Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Product Information**

Application	WB
Primary Accession	<a href="#">P31260</a>
Other Accession	<a href="#">NP_061824.3</a> , <a href="#">3206</a>
Reactivity	Human
Predicted	Human, Mouse, Pig, Cow
Host	Goat
Isotype	IgG
Calculated MW	42414

**Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Additional Information****Gene ID** 3206**Other Names**

Homeobox protein Hox-A10, Homeobox protein Hox-1.8, Homeobox protein Hox-1H, PL, HOXA10, HOX1H

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-HOXA10 Antibody (internal region), Biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Protein Information****Name** HOXA10**Synonyms** HOX1H**Function**

Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Binds to the DNA sequence 5'-AA[AT]TTTATTAC-3'.

**Cellular Location**


Nucleus.

**Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Images**



Biotinylated AF4299a (0.2 µg/ml) staining of Human Skeletal Muscle lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.