

Goat Anti-BMP7 Antibody (aa120-131) (internal region)
Catalog # AF4301a**Specification**

Goat Anti-BMP7 Antibody (aa120-131) (internal region) - Product Information

Application	WB
Primary Accession	P18075
Other Accession	NP_001710.1 , 85272 , 12162 , 655
Reactivity	Pig
Predicted	Human, Mouse, Rat, Pig, Cow, Dog
Host	Goat
Isotype	IgG
Calculated MW	49313

Goat Anti-BMP7 Antibody (aa120-131) (internal region) - Additional Information**Gene ID** 655**Other Names**

Bone morphogenetic protein 7, BMP-7, Osteogenic protein 1, OP-1, Eptoterminalfa, BMP7, OP1

Immunogen

This antibody is expected to recognize precursor only.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-BMP7 Antibody (aa120-131) (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-BMP7 Antibody (aa120-131) (internal region) - Protein Information**Name** BMP7**Synonyms** OP1**Function**

Growth factor of the TGF-beta superfamily that plays important role in various biological processes, including embryogenesis, hematopoiesis, neurogenesis and skeletal morphogenesis (PubMed: [31208997](http://www.uniprot.org/citations/31208997)). Initiates the canonical BMP signaling cascade by associating with type I receptor ACVR1 and type II receptor ACVR2A (PubMed: [9748228](http://www.uniprot.org/citations/9748228), PubMed: [12667445](http://www.uniprot.org/citations/12667445)). Once all three components are bound together in a complex at the cell surface, ACVR2A phosphorylates and activates ACVR1. In turn, ACVR1 propagates signal

by phosphorylating SMAD1/5/8 that travel to the nucleus and act as activators and repressors of transcription of target genes (PubMed:12478285). For specific functions such as growth cone collapse in developing spinal neurons and chemotaxis of monocytes, uses also BMPR2 as type II receptor (PubMed:31208997). Can also signal through non-canonical pathways such as P38 MAP kinase signaling cascade that promotes brown adipocyte differentiation through activation of target genes, including members of the SOX family of transcription factors (PubMed:27923061). Promotes the expression of HAMP, this is repressed by its interaction with ERFE (PubMed:30097509).

Cellular Location

Secreted.

Tissue Location

Expressed in the kidney and bladder. Lower levels seen in the brain

Goat Anti-BMP7 Antibody (aa120-131) (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-BMP7 Antibody (aa120-131) (internal region) - Images



AF4301a (1 µg/ml) staining of Pig Kidney lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.