

T Antibody (Ascites)

Mouse Monoclonal Antibody (Mab)
Catalog # AM1993a

Specification

T Antibody (Ascites) - Product Information

Application WB,E
Primary Accession O15178

Other Accession <u>Q07998</u>, <u>NP 003172.1</u>

Reactivity
Predicted
Predicted
Host
Clonality
Human
Zebrafish
Mouse
Monoclonal

Isotype IgM
Calculated MW 47443
Antigen Region 201-228

T Antibody (Ascites) - Additional Information

Gene ID 6862

Other Names

Brachyury protein, Protein T, T

Target/Specificity

This T antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 201-228 amino acids from human T.

Dilution

WB~~1:1000~8000

Format

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

T Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

T Antibody (Ascites) - Protein Information

Name TBXT (HGNC:11515)

Synonyms T



Function Involved in the transcriptional regulation of genes required for mesoderm formation and differentiation. Binds to a palindromic T site 5'-TTCACACCTAGGTGTGAA-3' DNA sequence and activates gene transcription when bound to such a site.

Cellular Location Nucleus

Tissue Location


Detected in testis, but not in other, normal tissues. Detected in lung tumors (at protein level)

T Antibody (Ascites) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

T Antibody (Ascites) - Images



T Antibody (Cat. #AM1993a) western blot analysis in NCI-H292 cell line lysates ($35\mu g$ /lane). This demonstrates the T antibody detected the T protein (arrow).

T Antibody (Ascites) - Background

The protein encoded by this gene is an embryonic nuclear transcription factor that binds to a specific DNA element, the palindromic T-site. It binds through a region in its N-terminus, called the T-box, and effects transcription of genes required for mesoderm formation and differentiation. The protein is localized to notochord-derived cells.

T Antibody (Ascites) - References

Jambhekar, N.A., et al. Arch. Pathol. Lab. Med. 134(8):1181-1187(2010)





Docherty, S.J., et al. BMC Genet. 11, 61 (2010): Yang, X.R., et al. Nat. Genet. 41(11):1176-1178(2009)
Oakley, G.J., et al. Mod. Pathol. 21(12):1461-1469(2008)
Bondue, A., et al. Cell Stem Cell 3(1):69-84(2008)