

Phospho-Ser603 Synapsin I Antibody

Affinity purified rabbit polyclonal antibody Catalog # AN1025

Specification

Phospho-Ser603 Synapsin I Antibody - Product Information

Application Primary Accession Reactivity Predicted Host Clonality Calculated MW WB <u>P17599</u> Rat Bovine, Human, Mouse, Xenopus, Zebrafish Rabbit polyclonal 78 KDa

Phospho-Ser603 Synapsin I Antibody - Additional Information

Gene ID Gene Name **Other Names** Synapsin-1, Synapsin I, SYN1

281510 SYN1

Target/Specificity

Synthetic phospho-peptide corresponding to amino acid residues surrounding Ser603 conjugated to KLH.

Dilution WB~~ 1:1000

Format Prepared from rabbit serum by affinity purification via sequential chromatography on phosphoand dephosphopeptide affinity columns.

Antibody Specificity

Specific for \sim 78k synapsin I doublet protein phosphorylated at Ser603.Immunolabeling of the synapsin I band is blocked by λ -phosphatase treatment.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-Ser603 Synapsin I Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice


Phospho-Ser603 Synapsin I Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Phospho-Ser603 Synapsin I Antibody - Images



Western blot of rat cortex lysate showing specific immunolabeling of the ~78k synapsin I phosphorylated at Ser603 (Control). The phosphospecificity of this labeling is shown in the second lane (lambda-phosphatase: λ -Ptase). The blot is identical to the control except that it was incubated in λ -Ptase (1200 units for 30 min) before being exposed to the phospho-Ser603 synapsin I antibody. The immunolabeling is completely eliminated by treatment with λ -Ptase.

Phospho-Ser603 Synapsin I Antibody - Background

Synapsin I plays a key role in synaptic plasticity in brain (Feng et al., 2002; Nayak et

al., 1996). This effect is due in large part to the ability of the synapsins to regulate the availability of synaptic vesicles for release. Th

e role of synapsin in synaptic plasticity and in synaptogensis is

regulated by phosphorylation (Jovanovic et al., 2001; Kao et al., 2002). Serine 603 is the site on synapsin I that is phosphorylated by calcium calmodulin kinase II and by p21-activated kinases (Sakurada et al., 2002; Czernik et al., 1987). Phosphorylation of this site is thought to regulate synaptic vesicle function (Nayak et al., 1996; Bahler and Greengard, 1987; McGuinness et al., 1989).

Phospho-Ser603 Synapsin I Antibody - References

Bahler M, Greengard P (1987) Synapsin I bundles F-actin
in a phosphorylation-dependent manner. Nature (London)
326:704-707.
Czernik AJ, Pang DT, Greengard P (1987) Amino acid
sequences surrounding the cAMP-dependent and
calcium/calmodulin-dependent phosphorylation sites in rat and bovine synapsin I. Proc Natl Acad
Sci (USA)
84:7518-7522.
Feng J, Chi P, Blanpied TA, Xu YM, Magarinos AM, Fe



rreira A, Takahashi RH, Kao HT, McEwen BS, Ryan TA, Augustine GJ, Greengard P (2002) Regulation of neurotransmitter release by synapsin III. J Neurosci 22:4372-4380. Jovanovic JN, Sihra TS, Nairn AC, Hemmings HC, Jr., Gr eengard P, Czernik AJ (2001) Opposing changes in phosphorylation of specific sites in synapsin I during Ca 2 +-dependent glutamate release in isolated nerve terminals. | Neurosci 21:7944-7953. Kao HT, Song HJ, Porton B, Ming GL, Hoh J, Abraham M, Czernik AJ, Pieribone VA, Poo MM, Greengard P (2002) A protein kinase A-dependent molecular switch in synapsin s regulates neurite outgrowth. Nature Neurosci 5:431-437. McGuinness TL, Brady ST, Gruner JA, Sugimori M, L linás RR, Greengard P (1989) Phosphorylation-dependent inhibition by synapsin I of organelle movement in squid axoplasm. J Neurosci 9:4138-4149. Nayak AS, Moore CI, Browning MD (1996) CAM kinase II phos phorylation of the presyn aptic protein synapsin is persistently increased during expression of long-term po tentiation. Proc Natl Acad Sci (USA) 93:15451-15456. Sakurada K, Kato H, Nagumo H, Hiraoka H, Furuya K, Ikuhar a T, Yamakita Y, Fukunaga K, Miyamoto E, Matsumura F, Matsuo YI, Naito Y, Sasaki Y (2002) Synapsin I is phosphorylated at Ser 603 by p21-activated kinases (PAKs) in vitro and in PC12 cells stimulated with bradykinin. | Biol Chem 277:45473-45479. Sergio Leal-Ortiz, Clarissa L. Waites, Ryan Terry-Lorenzo , Pedro Zamorano, Eckart D. Gundelfinger, and Craig C. Garner (2008) Piccolo modulation of Synapsin1a dynamics regulates synaptic vesicle exocytosis

J. Cell Biol., 181: 831 - 846.