

Phospho Ser862 mGluR7 Antibody

Affinity purified rabbit polyclonal antibody Catalog # AN1202

Specification

Phospho Ser862 mGluR7 Antibody - Product Information

Application FC, WB
Primary Accession P35400
Reactivity Mouse, Rat

Predicted Bovine, Human, Monkey, Zebrafish

Host Rabbit
Clonality polyclonal
Calculated MW 102 KDa

Phospho Ser862 mGluR7 Antibody - Additional Information

Gene ID 81672
Gene Name GRM7

Other Names

Metabotropic glutamate receptor 7, mGluR7, Grm7, Gprc1g, Mglur7

Target/Specificity

Synthetic phospho-peptide corresponding to amino acid residues surrounding Ser862 conjugated to KLH.

Dilution

FC~~1:500 WB~~ 1:1000

Format

Prepared from rabbit serum by affinity purification via sequential chromatography on phospho and dephospho-peptide columns.

Antibody Specificity

Specific for the $\sim 102 \text{k}$ mGluR7 protein phosphorylated at Ser862. Immunolabeling is blocked by preadsorption of antibody with the phospho-peptide used as antigen but not by the corresponding dephosphopeptide.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho Ser862 mGluR7 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping

Blue Ice




Phospho Ser862 mGluR7 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Phospho Ser862 mGluR7 Antibody - Images



Flow cytometric analysis of SH-SY5Y cells using Phospho Ser862 mGluR7 Antibody(green, Cat#AN1202) compared to an isotype control of rabbit IgG(blue). AN1202 was diluted at 1:500 dilution. An Alexa Fluor® 488 goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody.



Western blot of mouse brain lysate showing the specific immunolabeling the ~102k mGluR7 protein phosphorylated at Ser862. Immunolabeling blocked by the phospho-peptide used as antigen (peptide) but not by the corresponding dephospho-peptide (not shown).

Phospho Ser862 mGluR7 Antibody - Background

Metabotropic glutamate receptors (mGluRs) are key receptors in the modulation of excitatory





Tel: 858.875.1900 Fax: 858.875.1999

synaptic transmission in the central nervous system. They are implicated in many forms of neural plasticity as well as learning and memory and drug abuse (Bhattacharya et al., 2004; Francesconi et al., 2004; Wilson and Nicoll, 2001). The mGluRs are divided into three groups based on sequence identity and pharmacological properties: group I (mGluR1 and mGluR5) are localized in the perisynaptic region of the postsynaptic membrane, whereas group II (mGlur2 and mGluR3) and group III (mGluR4,6,7 and 8) are localized predominantly at presynaptic terminals. PKC phosphorylation of serine 862 on mGluR7 has been shown to be critical for stabilizing receptor surface expression and promoting binding to the synaptic PDZ-domain-containing protein PICK1 (Suh et al., 2008).

Phospho Ser862 mGluR7 Antibody - References

Bhattacharya M, Babwah AV, Godin C, Anborgh PH, Dale LB, Poulter MO, Ferguson SSG (2004) Ral and phospholipase D2-dependent pathway for constitutive metabotropic glutamate receptor endocytosis. J Neurosci 24:8752-8761.

Francesconi W, Cammalleri M, Sanna PP (2004) The metabotropic glutamate receptor 5 is necessary for late-phase long-term potentiation in the hippocampal CA1 region. Brain Res 1022:12-18.

Wilson RI, Nicoll RA (2001) Endogenous cannabinoids mediate retrograde signalling at hippocampal synapses. Nature (London) 410:588-592.

Suh YH, Pelkey KA, Lavezzari G, Roche PA, Huganir RL, McBain CJ, Roche KW. (2008) Corequirement of PICK1 binding and PKC phosphorylation for stable surface expression of the metabotropic glutamate receptor mGluR7. Neuron. 58(5):736-48