

Phospho-Ser239 ATRIP Antibody
Affinity purified rabbit polyclonal antibody
Catalog # AN1255

Specification

Phospho-Ser239 ATRIP Antibody - Product Information

Application	WB
Primary Accession	Q8WXE1
Reactivity	Mouse
Predicted	Bovine, Human, Monkey
Host	Rabbit
Clonality	polyclonal
Calculated MW	86 KDa

Phospho-Ser239 ATRIP Antibody - Additional Information

Gene ID	84126
Gene Name	ATRIP
Other Names	
ATR-interacting protein, ATM and Rad3-related-interacting protein, ATRIP, AGS1	

Target/Specificity

Synthetic phospho-peptide corresponding to amino acid residues surrounding Ser239 conjugated to KLH.

Dilution

WB~~ 1:500

Format

Prepared from rabbit serum by affinity purification via sequential chromatography on phospho- and dephospho-peptide affinity columns.

Antibody Specificity

Specific for the ~86k ATRIP protein phosphorylated at Ser239. The antibody also recognizes a band at ~32k.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-Ser239 ATRIP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Phospho-Ser239 ATRIP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-Ser239 ATRIP Antibody - Images



Western blot of mouse testis lysate showing specific immunolabeling of the ~86kd ATRIP protein phosphorylated at Ser239 (control). Phosphospecificity is shown in the second lane (lambda-phosphatase: λ-Ptase). The blot is identical to the control except that the lysate was incubated in λ-Ptase (800 units/1mg protein for 30 min). The immunolabeling is completely eliminated by treatment with λ-Ptase.

Phospho-Ser239 ATRIP Antibody - Background

ATRIP, ATR interacting protein, binds to ATR to regulate ATR expression, and is an essential component of the DNA damage checkpoint pathway (Cortez et al, 2001). ATR is recruited to DNA lesions in part through its association with ATRIP, which in turn interacts with the single-stranded DNA binding protein RPA (Ball et al, 2007). DNA replication forks may stall as a result of DNA damage causing phosphorylation of several proteins, including BRCA1 when colocalizing with ATR/ATRIP complex and RPA (Venere et al, 2007). The DNA replication fork stall coincides with BRCA1 directly phosphorylating ATRIP at ser239 (Venere et al, 2007).

Phospho-Ser239 ATRIP Antibody - References

- Cortez D, Guntuku S, Qin J, Elledge SJ (2001) ATR and ATRIP: partners in checkpoint signaling. *Science* 294(5547):1713-6.
- Ball HL, Ehrhardt MR, Mordes DA, Glick GG, Chazin WJ, Cortez D. (2007) Function of a conserved checkpoint recruitment domain in ATRIP proteins. *Mol Cell Biol.* 27(9):3367-77.
- Venere M, Synder A, Zgheib, and Halazonetis T. (2007) Phosphorylation of ATR-interacting Protein on Ser239 Mediates an Interaction with Breast-Ovarian Cancer Susceptibility 1 and Checkpoint Function. *Cancer Res* 67(13): 6100-5.