

C12orf30 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP10270C

Specification

C12orf30 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q14CX7
Other Accession	NP_079229.2
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	112292
Antigen Region	720-748

C12orf30 Antibody (Center) - Additional Information

Gene ID 80018

Other Names

N-alpha-acetyltransferase 25, NatB auxiliary subunit, Mitochondrial distribution and morphology protein 20, N-terminal acetyltransferase B complex subunit MDM20, NatB complex subunit MDM20, N-terminal acetyltransferase B complex subunit NAA25, p120, NAA25, C12orf30, MDM20, NAP1

Target/Specificity

This C12orf30 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 720-748 amino acids from the Central region of human C12orf30.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

C12orf30 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

C12orf30 Antibody (Center) - Protein Information

Name NAA25

Synonyms C12orf30, MDM20, NAP1

Function Non-catalytic subunit of the NatB complex which catalyzes acetylation of the N-terminal methionine residues of peptides beginning with Met-Asp, Met-Glu, Met-Asn and Met-Gln. May play a role in normal cell-cycle progression.

Cellular Location


Cytoplasm.

C12orf30 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C12orf30 Antibody (Center) - Images



Anti-C12orf30 Antibody (Center) at 1:1000 dilution + HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 112 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



C12orf30 Antibody (Center) (Cat. #AP10270c) western blot analysis in mouse cerebellum tissue lysates (15ug/lane). This demonstrates the C12orf30 antibody detected C12orf30 protein (arrow).

C12orf30 Antibody (Center) - Background

This gene encodes the auxiliary subunit of the heteromeric N-terminal acetyltransferase B complex. This complex acetylates methionine residues that are followed by acidic or asparagine residues.

C12orf30 Antibody (Center) - References

Feng, T., et al. Hum. Genet. 128(3):269-280(2010)
Pierce, B.L., et al. Hum. Hered. 69(3):193-201(2010)
Douroudis, K., et al. BMC Med. Genet. 11, 11 (2010) :
Pralhad, S., et al. Arthritis Rheum. 60(7):2124-2130(2009)
Polevoda, B., et al. BMC Proc 3 SUPPL 6, S2 (2009) :

C12orf30 Antibody (Center) - Citations

- [pTSara-NatB, an improved N-terminal acetylation system for recombinant protein expression in E. coli.](#)