

JMJD1B Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP1027A**Specification**

JMJD1B Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q7LBC6
Other Accession	Q6ZPY7
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	191581
Antigen Region	25-57

JMJD1B Antibody (N-term) - Additional Information**Gene ID** 51780**Other Names**

Lysine-specific demethylase 3B, 11411-, JmjC domain-containing histone demethylation protein 2B, Jumonji domain-containing protein 1B, Nuclear protein 5qNCA, KDM3B, C5orf7, JHDM2B, JMJD1B, KIAA1082

Target/Specificity

This JMJD1B antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 25-57 amino acids from the N-terminal region of human JMJD1B.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

JMJD1B Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

JMJD1B Antibody (N-term) - Protein Information**Name** KDM3B

Synonyms C5orf7, JHDM2B, JMJD1B, KIAA1082

Function Histone demethylase that specifically demethylates 'Lys-9' of histone H3, thereby playing a central role in histone code. Demethylation of Lys residue generates formaldehyde and succinate. May have tumor suppressor activity.

Cellular Location

Nucleus.

Tissue Location


Ubiquitous. Highly expressed in placenta, skeletal muscle, kidney, heart and liver.

JMJD1B Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

JMJD1B Antibody (N-term) - Images



Western blot analysis of anti-JMJD1B (N-term) Pab in Ramos cell line lysate. JMJD1B (N-term)(arrow) was detected using the purified Pab.

JMJD1B Antibody (N-term) - Background

Covalent modification of histones plays critical role in regulating chromatin structure and transcription. While most covalent histone modifications are reversible, only recently has it been established that methyl groups are subject to enzymatic removal from histones. A family of novel JmjC domain-containing histone demethylation (JHDM) enzymes have been identified that perform this specific function. Histone demethylation by JHDM proteins requires cofactors Fe(II) and alpha-ketoglutarate. Family members include JHDM1 (demethylating histone 3 at lysine 36), and JHDM2A as well as JMJD2CH3K9 (both of which demethylate histone 3 at lysine 9). Contributions of histone demethylase activity to tumor development, decreases in cell proliferation, and

hormone-dependent transcriptional activation have been observed.

JMJD1B Antibody (N-term) - References

- Katoh, M., et al., Int. J. Mol. Med. 12(5):817-821 (2003).
Hu, Z., et al., Oncogene 20(47):6946-6954 (2001).
Lai, F., et al., Genomics 71(2):235-245 (2001).
Lai, F., et al., Genomics 70(1):123-130 (2000).
Dias Neto, E., et al., Proc. Natl. Acad. Sci. U.S.A. 97(7):3491-3496 (2000).