

TNK1 Antibody (Center)

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP7722c

Specification

TNK1 Antibody (Center) - Product Information

Application WB, IHC-P,E Primary Accession Q13470

Other Accession Q99ML2, Q95364

Reactivity
Predicted
Host
Clonality
Polyclonal
Isotype
Antigen Region
Human
Mouse
Rabbit
Polyclonal
Rabbit IgG

TNK1 Antibody (Center) - Additional Information

Gene ID 8711

Other Names

Non-receptor tyrosine-protein kinase TNK1, CD38 negative kinase 1, TNK1 {ECO:0000312|EMBL:AAC994121}

Target/Specificity

This TNK1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 256-286 amino acids from the Central region of human TNK1.

Dilution

WB~~1:1000 IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TNK1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

TNK1 Antibody (Center) - Protein Information

Name TNK1 {ECO:0000312|EMBL:AAC99412.1}





Tel: 858.875.1900 Fax: 858.875.1999

Function Involved in negative regulation of cell growth. Has tumor suppressor properties. Plays a negative regulatory role in the Ras-MAPK pathway. May function in signaling pathways utilized broadly during fetal development and more selectively in adult tissues and in cells of the lymphohematopoietic system. Could specifically be involved in phospholipid signal transduction.

Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein

Tissue Location


Expressed in all umbilical cord blood, bone marrow and adult blood cell sub-populations and in several leukemia cell lines. Highly expressed in fetal blood, brain, lung, liver and kidney Detected at lower levels in adult prostate, testis, ovary, small intestine and colon. Not expressed in adult lung, liver, kidney or brain.

TNK1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

TNK1 Antibody (Center) - Images



Western blot analysis of anti-TNK1 pab (cat# AP7722c) in Hela cell line lysate. Dilution of anti-TNK1 was 1:100; dilution of secondary antibody (goat anti-rabbit-HRP) was 1:7000. Data and protocol courtesy of Dr. Richard Lu, Partners HealthCare System at Harvard University.





Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

TNK1 Antibody (Center) - Background

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the g phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains.

TNK1 Antibody (Center) - References

Blume-Jensen P, et al. Nature 2001. 411: 355. Cantrell D, J. Cell Sci. 2001. 114: 1439. Jhiang S Oncogene 2000. 19: 5590. Manning G, et al. Science 2002. 298: 1912. Moller, D, et al. Am. J. Physiol. 1994. 266: C351-C359.

Robertson, S. et al. Trends Genet. 2000. 16: 368.

Robinson D, et al. Oncogene 2000. 19: 5548.

Van der Ven, P, et al. Hum. Molec. Genet. 1993. 2: 1889.

Vanhaesebroeck, B, et al. Biochem. J. 2000. 346: 561.

Van Weering D, et al. Recent Results Cancer Res. 1998. 154: 271.

TNK1 Antibody (Center) - Citations

 High-throughput RNAi screening identifies a role for TNK1 in growth and survival of pancreatic cancer cells.