

Anti-Goat IgG F(c) (Rhodamine Conjugated) Secondary Antibody

Rabbit Polyclonal, Rhodamine (TRITC)
Catalog # ASR1766

Specification

Target Species

Physical State

Immunogen

Anti-Goat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Product Information

Description Anti-GOAT IgG F(c) (RABBIT) Antibody

Rhodamine Conjugated

Host Rabbit

Conjugate Rhodamine (TRITC)

FP Value 4.0 moles Rhodamine (TRITC) per mole of

lgG Goat Polyclonal

Clonality Polyce Application ,3,4,

Application Note FLISA 1:10,000-1:50,000;IF Microscopy

1:1,000-1:5,000;FlowCytometry

1:500-1:2,500 Lyophilized

Host Isotype IgG
Target Isotype IgG F(c)

Buffer 0.02 M Potassium Phosphate, 0.15 M

Sodium Chloride, pH 7.2 Goat IgG F(c) fragment

Reconstitution Volume 1.0 mL

Reconstitution Buffer Restore with deionized water (or

equivalent)

Stabilizer 10 mg/mL Bovine Serum Albumin (BSA) -

Immunoglobulin and Protease free

Preservative 0.01% (w/v) Sodium Azide

Anti-Goat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Additional Information

Shipping Condition

Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Goat IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Rabbit Serum, Goat IgG. Goat IgG F(c), and Goat Serum. No reaction was observed against Goat IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Goat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protein Information

Anti-Goat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Goat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Images

Anti-Goat IgG F(c) (Rhodamine Conjugated) Secondary Antibody - Background

This product is designed for immunofluorescence microscopy, fluorescence based plate assays (FLISA) and fluorescent western blotting. This product is also suitable for multiplex analysis, including multicolor imaging, utilizing various commercial platforms.