

# Anti-Ferret IgM (mu chain) (Biotin Conjugated) Secondary Antibody

Goat Polyclonal, Biotin Catalog # ASR2122

## Specification

Description

# Anti-Ferret IgM (mu chain) (Biotin Conjugated) Secondary Antibody - Product Information

Host Conjugate Target Species Clonality Application Application Note

Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer

Preservative

Anti-FERRET IgM (mu chain) (GOAT) **Antibody Biotin Conjugated** Goat **Biotin** Ferret Polyclonal ,1,10,15, ELISA 1:5,000-1:20,000;Western Blot 1:2,000-1:10,000;Immunochemistry 1:1,000-1:5,000 Lyophilized laG IgM μ chain 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 Ferret IgM mu heavy chain 1.0 mL Restore with deionized water (or equivalent) 10 mg/mL Bovine Serum Albumin (BSA) -**Immunoglobulin and Protease free** 0.01% (w/v) Sodium Azide

# Anti-Ferret IgM (mu chain) (Biotin Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

#### **Purity**

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Ferret IgM coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-biotin, anti-Goat Serum, Ferret IgM and Ferret Serum. No reaction was observed against Ferret IgA or Ferret IgG.

### Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

**Precautions Note** 



This product is for research use only and is not intended for therapeutic or diagnostic applications.

Anti-Ferret IgM (mu chain) (Biotin Conjugated) Secondary Antibody - Protein Information

## Anti-Ferret IgM (mu chain) (Biotin Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# Anti-Ferret IgM (mu chain) (Biotin Conjugated) Secondary Antibody - Images