

Anti-Bovine IgG F(c) (Peroxidase Conjugated) Secondary Antibody Rabbit Polyclonal, Peroxidase (Horseradish)

Catalog # ASR2648

Specification

Anti-Bovine IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Product Information

Description

- Host Conjugate Target Species Clonality Application Application Note
- Physical State Host Isotype Target Isotype Buffer

Immunogen Reconstitution Volume Reconstitution Buffer

Stabilizer Preservative Anti-BOVINE IgG F(c) (RABBIT) Antibody **Peroxidase Conjugated** Rabbit Peroxidase (Horseradish) **Bovine** Polyclonal ,1,10,15, ELISA 1:10,000-1:50,000;Western Blot 1:1,000-1:10,000;Immunochemistry 1:500-1:2,500 Lyophilized laG laG F(c) 0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2 **Bovine IgG F(c) fragment** 1.0 mL Restore with deionized water (or equivalent) 10 mg/ml Polvethvlene Glvcol (PEG-8000) 0.01% (w/v) Gentamicin Sulfate. Do NOT add Sodium Azide!

Anti-Bovine IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Additional Information

Shipping Condition Ambient

Purity

This product was prepared from monospecific antiserum by immunoaffinity chromatography using Bovine IgG coupled to agarose beads. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Peroxidase, anti-Rabbit Serum, Bovine IgG, Bovine IgG F(c) and Bovine Serum. No reaction was observed against Bovine IgG F(ab')2.

Storage Condition

Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.

Precautions Note

This product is for research use only and is not intended for therapeutic or diagnostic applications.



Anti-Bovine IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protein Information

Anti-Bovine IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Bovine IgG F(c) (Peroxidase Conjugated) Secondary Antibody - Images