

AKR1C2 Antibody (monoclonal) (M03)

Mouse monoclonal antibody raised against a partial recombinant AKR1C2. Catalog # AT1093a

Specification

AKR1C2 Antibody (monoclonal) (M03) - Product Information

Application WB, E **Primary Accession** P52895 Other Accession BC063574 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a Kappa Calculated MW 36735

AKR1C2 Antibody (monoclonal) (M03) - Additional Information

Gene ID 1646

Other Names

Aldo-keto reductase family 1 member C2, 1---, 3-alpha-HSD3, Chlordecone reductase homolog HAKRD, Dihydrodiol dehydrogenase 2, DD-2, DD2, Dihydrodiol dehydrogenase/bile acid-binding protein, DD/BABP, Trans-1, 2-dihydrobenzene-1, 2-diol dehydrogenase, Type III 3-alpha-hydroxysteroid dehydrogenase, AKR1C2, DDH2

Target/Specificity

AKR1C2 (AAH63574, 224 a.a. ~ 323 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

AKR1C2 Antibody (monoclonal) (M03) is for research use only and not for use in diagnostic or therapeutic procedures.

AKR1C2 Antibody (monoclonal) (M03) - Protocols


Provided below are standard protocols that you may find useful for product applications.

Western Blot




- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

AKR1C2 Antibody (monoclonal) (M03) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.74 KDa) .




Western Blot analysis of AKR1C2 expression in transfected 293T cell line by AKR1C2 monoclonal antibody (M03), clone 3C11.

Lane 1: AKR1C2 transfected lysate(36.7 KDa).

Lane 2: Non-transfected lysate.





Detection limit for recombinant GST tagged AKR1C2 is approximately 0.3ng/ml as a capture antibody.

AKR1C2 Antibody (monoclonal) (M03) - Background

This gene encodes a member of the aldo/keto reductase superfamily, which consists of more than 40 known enzymes and proteins. These enzymes catalyze the conversion of aldehydes and ketones to their corresponding alcohols using NADH and/or NADPH as cofactors. The enzymes display overlapping but distinct substrate specificity. This enzyme binds bile acid with high affinity, and shows minimal 3-alpha-hydroxysteroid dehydrogenase activity. This gene shares high sequence identity with three other gene members and is clustered with those three genes at chromosome 10p15-p14.

AKR1C2 Antibody (monoclonal) (M03) - References

1.Aldo-keto reductases AKR1C1, AKR1C2 and AKR1C3 may enhance progesterone metabolism in ovarian endometriosis. Hevir N, Vouk K, Sinkovec J, Ribic-Pucelj M, Lanisnik Rizner T.Chem Biol Interact. 2011 Jan 11. [Epub ahead of print]2.Aldo-keto reductase 1C2 fails to metabolize doxorubicin and daunorubicin in vitro. Takahashi RH, Bains OS, Pfeifer TA, Grigliatti TA, Reid RE, Riggs KW. Drug Metab Dispos. 2008 Jun; 36(6):991-4. Epub 2008 Mar 5.