

ARHGEF1 Antibody (monoclonal) (M02)

Mouse monoclonal antibody raised against a partial recombinant ARHGEF1. Catalog # AT1184a

Specification

ARHGEF1 Antibody (monoclonal) (M02) - Product Information

Application WB, E **Primary Accession** 092888 Other Accession BC034013 Reactivity Human Host mouse Clonality **Monoclonal** Isotype IgG2a Kappa Calculated MW 102435

ARHGEF1 Antibody (monoclonal) (M02) - Additional Information

Gene ID 9138

Other Names

Rho guanine nucleotide exchange factor 1, 115 kDa guanine nucleotide exchange factor, p115-RhoGEF, p115RhoGEF, Sub15, ARHGEF1

Target/Specificity

ARHGEF1 (AAH34013.2, 830 a.a. \sim 927 a.a) partial recombinant protein with GST tag. MW of the GST tag alone is 26 KDa.

Dilution

WB~~1:500~1000

Format

Clear, colorless solution in phosphate buffered saline, pH 7.2.

Storage

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Precautions

ARHGEF1 Antibody (monoclonal) (M02) is for research use only and not for use in diagnostic or therapeutic procedures.

ARHGEF1 Antibody (monoclonal) (M02) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot




- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ARHGEF1 Antibody (monoclonal) (M02) - Images



Antibody Reactive Against Recombinant Protein. Western Blot detection against Immunogen (36.41 KDa).




Western Blot analysis of ARHGEF1 expression in transfected 293T cell line by ARHGEF1 monoclonal antibody (M02), clone 2D2.

Lane 1: ARHGEF1 transfected lysate (Predicted MW: 102.4 KDa).

Lane 2: Non-transfected lysate.





Detection limit for recombinant GST tagged ARHGEF1 is approximately 1ng/ml as a capture antibody.

ARHGEF1 Antibody (monoclonal) (M02) - Background

Rho GTPases play a fundamental role in numerous cellular processes that are initiated by extracellular stimuli that work through G protein coupled receptors. The encoded protein may form complex with G proteins and stimulate Rho-dependent signals. Multiple alternatively spliced transcript variants have been found for this gene, but the full-length nature of some variants has not been defined.

ARHGEF1 Antibody (monoclonal) (M02) - References

Variation at the NFATC2 Locus Increases the Risk of Thiazolinedinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.Mutation of ARHGAP9 in patients with coronary spastic angina. Takefuji M, et al. J Hum Genet, 2010 Jan. PMID 19911011.P115 RhoGEF and microtubules decide the direction apoptotic cells extrude from an epithelium. Slattum G, et al. J Cell Biol, 2009 Sep 7. PMID 19720875.Reversible translocation of p115-RhoGEF by G(12/13)-coupled receptors. Meyer BH, et al. J Cell Biochem, 2008 Aug 1. PMID 18320579.