

DAZ2 Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP12295b**Specification**

DAZ2 Antibody (C-term) Blocking peptide - Product Information

Primary Accession [Q13117](#)

DAZ2 Antibody (C-term) Blocking peptide - Additional Information

Gene ID 57055

Other Names

Deleted in azoospermia protein 2, DAZ2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

DAZ2 Antibody (C-term) Blocking peptide - Protein Information

Name DAZ2

Function

RNA-binding protein that plays an essential role in spermatogenesis. May act by binding to the 3'-UTR of mRNAs and regulating their translation.

Cellular Location

Cytoplasm. Nucleus. Note=Predominantly cytoplasmic. Nuclear at some stages of spermatozoide development. Localizes both to the nuclei and cytoplasm of spermatozoide differentiation. Nuclear in fetal gonocytes and in spermatogonial nuclei. It then relocates to the cytoplasm during male meiosis

Tissue Location

Testis specific..

DAZ2 Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

DAZ2 Antibody (C-term) Blocking peptide - Images

DAZ2 Antibody (C-term) Blocking peptide - Background

This gene is a member of the DAZ gene family and is a candidate for the human Y-chromosomal azoospermia factor (AZF). Its expression is restricted to premeiotic germ cells, particularly in spermatogonia. It encodes an RNA-binding protein that is important for spermatogenesis. Four copies of this gene are found on chromosome Y within palindromic duplications; one pair of genes is part of the P2 palindrome and the second pair is part of the P1 palindrome. Each gene contains a 2.4 kb repeat including a 72-bp exon, called the DAZ repeat; the number of DAZ repeats is variable and there are several variations in the sequence of the DAZ repeat. Each copy of the gene also contains a 10.8 kb region that may be amplified; this region includes five exons that encode an RNA recognition motif (RRM) domain. This gene contains one copy of the 10.8 kb repeat. Alternative splicing results in multiple transcript variants encoding different isoforms.

DAZ2 Antibody (C-term) Blocking peptide - References

Kim, B., et al. Hum. Reprod. 24(6):1507-1515(2009) Lardone, M.C., et al. Fertil. Steril. 88(5):1318-1326(2007) A, Z.C., et al. Yi Chuan 28(9):1057-1060(2006) Zhang, F., et al. Ann. Hum. Genet. 70 (PT 3), 304-313 (2006) :Yang, Y., et al. Fertil. Steril. 85(4):1061-1063(2006)