

PION Antibody (C-term) Blocking peptide
Synthetic peptide
Catalog # BP12480b**Specification**

PION Antibody (C-term) Blocking peptide - Product InformationPrimary Accession [A4D1B5](#)**PION Antibody (C-term) Blocking peptide - Additional Information****Gene ID** 54103**Other Names**

Gamma-secretase-activating protein, GSAP, Protein pigeon homolog, Gamma-secretase-activating protein 16 kDa C-terminal form, GSAP-16K, GSAP, PION

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PION Antibody (C-term) Blocking peptide - Protein Information**Name** GSAP**Synonyms** PION**Function**

Regulator of gamma-secretase activity, which specifically activates the production of amyloid-beta protein (amyloid-beta protein 40 and amyloid-beta protein 42), without affecting the cleavage of other gamma-secretase targets such as Notch. The gamma-secretase complex is an endoprotease complex that catalyzes the intramembrane cleavage of integral membrane proteins such as Notch receptors and APP (amyloid-beta precursor protein). Specifically promotes the gamma- cleavage of APP CTF-alpha (also named APP-CTF) by the gamma-secretase complex to generate amyloid-beta, while it reduces the epsilon-cleavage of APP CTF-alpha, leading to a low production of AICD.

Cellular Location

Golgi apparatus, trans-Golgi network

Tissue Location

Widely expressed..

PION Antibody (C-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

PION Antibody (C-term) Blocking peptide - Images**PION Antibody (C-term) Blocking peptide - Background**

Accumulation of neurotoxic amyloid-beta is a major hallmark of Alzheimer disease (AD; MIM 104300). Formation of amyloid-beta is catalyzed by gamma-secretase (see PSEN1; MIM104311), a protease with numerous substrates. PION, or GSAP, selectively increases amyloid-beta production through a mechanism involving its interaction with both gamma-secretase and its substrate, the amyloid-beta precursor protein (APP; MIM 104760) C-terminal fragment (APP-CTF) (He et al., 2010 [PubMed20811458]).

PION Antibody (C-term) Blocking peptide - References

He, G., et al. Nature 467(7311):95-98(2010) Oh, J.H., et al. Mamm. Genome 16(12):942-954(2005)