

**CD48 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP8664b****Specification**

---

**CD48 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [P09326](#)**CD48 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 962**Other Names**

CD48 antigen, B-lymphocyte activation marker BLAST-1, BCM1 surface antigen, Leukocyte antigen MEM-102, SLAM family member 2, SLAMF2, Signaling lymphocytic activation molecule 2, TCT1, CD48, CD48, BCM1, BLAST1

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP8664b](/products/AP8664b) was selected from the C-term region of human CD48. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**CD48 Antibody (C-term) Blocking Peptide - Protein Information****Name** CD48**Synonyms** BCM1, BLAST1**Function**

Glycosylphosphatidylinositol (GPI)-anchored cell surface glycoprotein that interacts via its N-terminal immunoglobulin domain with cell surface receptors including 2B4/CD244 or CD2 to regulate immune cell function and activation (PubMed: [27249817](http://www.uniprot.org/citations/27249817), PubMed: [12007789](http://www.uniprot.org/citations/12007789)). Participates in T-cell signaling transduction by associating with CD2 and efficiently bringing the Src family protein kinase LCK and LAT to the TCR/CD3 complex (PubMed: [19494291](http://www.uniprot.org/citations/19494291)). In turn,

promotes LCK phosphorylation and subsequent activation (PubMed:<a href="http://www.uniprot.org/citations/12007789" target="\_blank">12007789</a>). Induces the phosphorylation of the cytoplasmic immunoreceptortyrosine switch motifs (ITSMs) of CD244 initiating a series of signaling events that leads to the generation of the immunological synapse and the directed release of cytolytic granules containing perforin and granzymes by T-lymphocytes and NK-cells (PubMed:<a href="http://www.uniprot.org/citations/9841922" target="\_blank">9841922</a>, PubMed:<a href="http://www.uniprot.org/citations/27249817" target="\_blank">27249817</a>).

**Cellular Location**

Cell membrane; Lipid-anchor, GPI-anchor. Secreted

**Tissue Location**

Widely expressed on all hematopoietic cells.

**CD48 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**CD48 Antibody (C-term) Blocking Peptide - Images****CD48 Antibody (C-term) Blocking Peptide - Background**

BLAST1 is the designation used for an activation-associated cell surface glycoprotein of 40 to 45 kD expressed primarily in mitogen-stimulated human lymphocytes. The protein sequence predicted by the cDNA encoding BLAST1 indicates that BLAST1 is a member of the immunoglobulin supergene family. Yokoyama (1991) identified the BLAST1 activation/adhesion molecule as CD48.

**CD48 Antibody (C-term) Blocking Peptide - References**

Ramos-Lopez,E., et.al., Tissue Antigens 68 (2), 147-152 (2006)Khan,N.A., et.al., Cell. Microbiol. 9 (1), 169-178 (2007)