

UNC13B Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP8905c**Specification**

UNC13B Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [O14795](#)**UNC13B Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 10497**Other Names**

Protein unc-13 homolog B, Munc13-2, munc13, UNC13B, UNC13

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP8905c](/products/AP8905c) was selected from the Center region of human UNC13B. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

UNC13B Antibody (Center) Blocking Peptide - Protein Information**Name** UNC13B ([HGNC:12566](#))**Synonyms** UNC13**Function**

Plays a role in vesicle maturation during exocytosis as a target of the diacylglycerol second messenger pathway. Is involved in neurotransmitter release by acting in synaptic vesicle priming prior to vesicle fusion and participates in the activity-depending refilling of readily releasable vesicle pool (RRP) (By similarity). Essential for synaptic vesicle maturation in a subset of excitatory/glutamatergic but not inhibitory/GABA-mediated synapses (By similarity). In collaboration with UNC13A, facilitates neuronal dense core vesicles fusion as well as controls the location and efficiency of their synaptic release (By similarity).

Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein. Cell membrane Synapse. Note=Localized to

synapses. Translocated to the plasma membrane in response to phorbol ester binding (By similarity)

Tissue Location

Expressed in kidney cortical epithelial cells and brain.

UNC13B Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

UNC13B Antibody (Center) Blocking Peptide - Images**UNC13B Antibody (Center) Blocking Peptide - Background**

UNC13B plays a role in vesicle maturation during exocytosis as a target of the diacylglycerol second messenger pathway. It is involved in neurotransmitter release by acting in synaptic vesicle priming prior to vesicle fusion and participates in the activity-dependent refilling of readily releasable vesicle pool (RRP). It is essential for synaptic vesicle maturation in a subset of excitatory/glutamatergic but not inhibitory/GABA-mediated synapses (By similarity).

UNC13B Antibody (Center) Blocking Peptide - References

Sjoeblom T., et.al., Science 314:268-274(2006). Yu L.-R., Zhu Z., et.al., J. Proteome Res. 6:4150-4162(2007).