

IL-20, human recombinant proteinIL10, D ZCYTO10, IL-20, MGC96907, Four α helix cytokine, Zcyto10,
Catalog # PBV10302r**Specification**

IL-20, human recombinant protein - Product infoPrimary Accession [O9NYY1](#)
Calculated MW **17.6 kDa KDa****IL-20, human recombinant protein - Additional Info**Gene ID **50604**
Gene Symbol **IL20**
Other Names
IL10, D ZCYTO10, IL-20, MGC96907, Four α helix cytokine, Zcyto10,
Gene Source **Human**
Source **E. coli**
Assay&Purity **SDS-PAGE; \geq 97%**
Assay2&Purity2 **HPLC; \geq 97%**
Recombinant **Yes**
Target/Specificity
IL-20**Application Notes**Reconstitute in H₂O to a concentration of 0.1 to 1.0 μ g/ μ l. It is recommended that further dilutions be made into buffer containing carrier protein or medium containing serum.**Format**

Lyophilized protein

Storage

-20°C; Sterile filtered and lyophilized from 10 mM Sodium Citrate, pH 3.0 + 100 mM NaCl.

IL-20, human recombinant protein - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IL-20, human recombinant protein - Images

IL-20, human recombinant protein - Background

Interleukin-20 has a 28% amino acid identity with IL-10 and induces keratinocyte proliferation. Recombinant human IL-20 is a 17.6 kDa protein consisting of 153 amino acids residues.

IL-20, human recombinant protein - References

Blumberg H., et al. Cell 104:9-19(2001).
Hsieh M.Y., et al. Genes Immun. 7:234-242(2006).
Clark H.F., et al. Genome Res. 13:2265-2270(2003).
Gregory S.G., et al. Nature 441:315-321(2006).
Zhang Z., et al. Protein Sci. 13:2819-2824(2004).