

# PDL1 Monoclonal Antibody

Purified Mouse Monoclonal Antibody (Mab)

**Catalog # AW5698**

Catalog # AW5698

Application: WB, IHC-P,E

Primary Accession: Q9NZQ7

Reactivity: Human


Predicted: Human

Host: Mouse


Clonality: Monoclonal

Isotype: IgG1


Antigen Source: Human




All lanes : Anti-PDL-1 Antibody at 0.5-1µg/ml dilution Lane 1: 293T/PDL1 whole cell lysate Lane 2: 293T whole cell lysate Lane 3: NCI-H1975 whole cell lysate Lane 4: MDA-MB231 whole cell lysate Lane 5: MCF-7 whole cell lysate Lane 6: NCI-H226 whole cell lysate Lysates/proteins at 30 µg per lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated at 1/5000 dilution. Predicted band size : 32 kDa Blocking/Dilution buffer: 5% NFDm/TBST.




Immunohistochemical analysis of PDL-1 in human non-small cell lung carcinoma sections (IHC-P, paraformaldehyde-fixed, paraffin-embedded sections) by Dako test. Tissue was fixed with formaldehyde; antigen retrieval was by heat mediation with EDTA buffer (pH9.0). Samples were incubated with primary antibody (0.85µg/ml) for 1 hour at room temperature. Aundiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Immunohistochemical analysis of PDL-1 in human non-small cell lung carcinoma sections (IHC-P, paraformaldehyde-fixed, paraffin-embedded sections) by Leica test. Tissue was fixed with formaldehyde; antigen retrieval was by heat mediation with EDTA buffer (pH9.0). Samples were incubated with primary antibody (0.85µg/ml) for 1 hour at room temperature. Aundiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Immunohistochemical analysis of PDL-1 in human non-small cell lung carcinoma sections (IHC-P, paraformaldehyde-fixed, paraffin-embedded sections) by Leica test. Tissue was fixed with formaldehyde; antigen retrieval was by heat mediation with EDTA buffer (pH9.0). Samples were incubated with primary antibody (0.85µg/ml) for 1 hour at room temperature. Aundiluted biotinylated goat polyvalent antibody was used as the secondary antibody.




**Inquire For Your Free Sample**  
[www.abcepta.com/freesample](http://www.abcepta.com/freesample)


# PDL1 Monoclonal Antibody

Purified Mouse Monoclonal Antibody (Mab)


Catalog # AW5698



Immunohistochemical analysis of PDL-1 in human tonsil tissue sections(IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde; antigen retrieval was by heat mediation with EDTA buffer (pH9.0). Samples were incubated with primary antibody (0.85µg/ml) for 1 hour at room temperature. Aundiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Immunohistochemical analysis of PDL-1 in human tonsil tissue sections(IHC-P - paraformaldehyde-fixed, paraffin-embedded sections) by Dako test. Tissue was fixed with formaldehyde; antigen retrieval was by heat mediation with EDTA buffer (pH9.0). Samples were incubated with primary antibody (0.85µg/ml) for 1 hour at room temperature. Aundiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Immunohistochemical analysis of PDL-1 in MCF-7 cell (left) and NCI-H226(right) cell sections . Cells were fixed with formaldehyde and blocked with super block for 10 minutes at room temperature; antigen retrieval was by heat mediation with EDTA buffer (pH9.0). Samples were incubated with primary antibody (0.85µg/ml) for 1 hour at room temperature. Aundiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

## Custom Services

- **20,000+** antibody projects completed
- **30,000+** peptides synthesized
- **WB, IHC, IF** validation available
- AAALAC and ISO accredited facility

✓ **Get Your Free Quote Today!**

Email: [sales@abcepta.com](mailto:sales@abcepta.com) or Tel: 888.735.7227



## Catalog Products

### PRIMARY ANTIBODIES (55,000+)

- |                         |                       |
|-------------------------|-----------------------|
| • Cancer                | • Metabolism          |
| • Cardiovascular        | • Microbiology        |
| • Cell Biology          | • Neuroscience        |
| • Developmental Biology | • Signal Transduction |
| • Immunology            | • Stem Cells          |

**SECONDARY ANTIBODIES**  
**CELL/TISSUE/LYSATES**  
**PEPTIDES**  
**FL cDNA CLONES**