

Anti-Human BMP-2 Antibody
Catalog # ABG10035**Specification****Anti-Human BMP-2 Antibody - Product Information**

Application	WB, IHC, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Anti-Human BMP-2 Antibody - Additional Information**Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hBMP-2. Anti-Human BMP-2 specific antibody was purified by affinity chromatography employing immobilized hBMP-2 matrix.

WesternBlot

To detect hBMP-2 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hBMP-2 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hBMP-2 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human BMP-2 (60-082BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hBMP-2.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of human colon/rectum adenocarcinoma. The recommended concentration is 0.25 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

 Additional Immunostaining data available. Please contact Tech Support for information.

Neutralization

To yield one-half maximal inhibition [**ND**50] of the biological activity of hBMP-2 (200 ng/ml), a concentration of 1.3-1.9 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human BMP-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human BMP-2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human BMP-2 Antibody - Images