

# **Anti-Human BMP-4 Antibody**

Catalog # ABG10037

# **Specification**

# **Anti-Human BMP-4 Antibody - Product Information**

Application WB, IHC, E
Reactivity Human
Host Mouse
Clonality Monoclonal

## **Anti-Human BMP-4 Antibody - Additional Information**

#### **Preparation**

Produced in mice using highly pure (>98%) recombinant human BMP-4 as the immunizing antigen. This IgG2<sub>bK</sub> antibody was purified from cell culture by Protein G affinity chromatography.

#### WesternBlot

To detect hBMP-4 by Western Blot analysis this antibody can be used at a concentration of 0.25- $0.50 \mu g/ml$ . Used in conjunction with compatible secondary reagents the detection limit for recombinant hBMP-4 is 2.0- $4.0 \mu g/lane$ , under non-reducing conditions.

### Sandwich

In a sandwich ELISA (assuming  $100\mu$ l/well), a concentration of 4.0- $8.0~\mu$ g/ml of this antibody will detect at least 2000 pg/ml of recombinant human BMP-4 when used in conjunction with compatible secondary reagents.

## **Immunohistochemistry**

This antibody stained HepG2 cells. The primary antibody was incubated at 3.0  $\mu$ g/ml overnight at 4°C followed by a fluorescent labeled secondary antibody. Optimal concentrations and conditions may vary. Information courtesy of the Cell Profiling group, SciLifeLab Stockholm.

#### **Neutralization**

To yield one-half maximal inhibition <strong>[ND<sub>50</sub>]</strong> of the biological activity of Human BMP-4 (5.0 ng/ml), a concentration of 0.4-0.6 µg/ml of this antibody is required.

#### Formulation

A sterile filtered antibody solution was lyophilized from PBS.

### Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

# **Storage**

-20°C

#### **Precautions**

Anti-Human BMP-4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



# **Anti-Human BMP-4 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

**Anti-Human BMP-4 Antibody - Images**