

Anti-Human CXCL16 Antibody
Catalog # ABG10057**Specification**

Anti-Human CXCL16 Antibody - Product Information

Application	WB, IHC, E
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Anti-Human CXCL16 Antibody - Additional Information**Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hCXCL16. Anti-Human CXCL16 specific antibody was purified by affinity chromatography employing immobilized hCXCL16 matrix.

WesternBlot

To detect hCXCL16 by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hCXCL16 is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hCXCL16 by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human CXCL16 (60-109BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hCXCL16.

Immunohistochemistry

This antibody stained formalin-fixed paraffin-embedded sections of human colorectal adenocarcinoma tissue. The recommended concentration is 0.5 µg/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Antigen retrieval is not required for these conditions. Optimal concentrations and conditions may vary.
*Additional Immunostaining data available. Please contact Tech Support for information.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human CXCL16 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human CXCL16 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Human CXCL16 Antibody - Images