

Anti-Murine Eotaxin Antibody

Catalog # ABG10082

Specification

Anti-Murine Eotaxin Antibody - Product Information

Application WB, E
Reactivity Mouse
Host Rabbit
Clonality Polyclonal

Anti-Murine Eotaxin Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mEotaxin. Anti-Murine Eotaxin specific antibody was purified by affinity chromatography employing immobilized mEotaxin matrix.

WesternBlot

To detect mEotaxin by Western Blot analysis this antibody can be used at a concentration of $0.1\text{-}0.2~\mu\text{g/ml}$. Used in conjunction with compatible secondary reagents the detection limit for recombinant mEotaxin is 1.5-3.0~ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect mEotaxin by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Murine Eotaxin (61-064BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mEotaxin.

Neutralization

To yield one-half maximal inhibition [ND₅₀] of the biological activity of mEotaxin (100.00 ng/ml), a concentration of 5.0-8.0 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.


Storage


-20°C

Precautions

Anti-Murine Eotaxin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Murine Eotaxin Antibody - Protocols







Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Murine Eotaxin Antibody - Images