

**Anti-Human GM-CSF Antibody**  
**Catalog # ABG10139****Specification**

---

**Anti-Human GM-CSF Antibody - Product Information**

Application	<b>WB, IHC, E</b>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>

**Anti-Human GM-CSF Antibody - Additional Information****Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hGM-CSF. Anti-Human GM-CSF specific antibody was purified by affinity chromatography employing immobilized hGM-CSF matrix.

**WesternBlot**

To detect hGM-CSF by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hGM-CSF is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

**Sandwich**

To detect hGM-CSF by sandwich ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Human GM-CSF (60-167BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hGM-CSF.

**Immunohistochemistry**

This antibody stained formalin-fixed, paraffin-embedded sections of human cervical squamous cell carcinoma. The recommended concentration is 0.25 µg/mL with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

**Neutralization**

To yield one-half maximal inhibition [**ND**"><sub>50</sub></span>] of the biological activity of hGM-CSF (1.0 ng/ml), a concentration of 0.13-0.20 µg/ml of this antibody is required.

**Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

**Reconstitution**

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

**Storage**

-20°C

**Precautions**

Anti-Human GM-CSF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### **Anti-Human GM-CSF Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Anti-Human GM-CSF Antibody - Images**