

# Anti-Human GRO-B Antibody

Catalog # ABG10147

# **Specification**

# Anti-Human GRO-β Antibody - Product Information

Application WB, IHC, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

## Anti-Human GRO-β Antibody - Additional Information

#### **Preparation**

Produced from sera of rabbits pre-immunized with highly pure recombinant Human GRO- $\beta$ . Anti-Human GRO- $\beta$  specific antibody was purified by affinity chromatography employing immobilized Human GRO- $\beta$  matrix.

#### WesternBlot

To detect human GRO- $\beta$  by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 µg/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Human GRO- $\beta$  is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

## Sandwich

To detect Human GRO- $\beta$  by sandwich ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.5 - 2.0  $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human GRO- $\beta$  (60-171BT) as a detection antibody, allows the detection of at least 0.2 - 0.4  $\mu$ g/well of recombinant Human GRO- $\beta$ .

### **Immunohistochemistry**

This antibody stained formalin-fixed paraffin-embedded sections of human colon/rectum adenocarcinoma tissue. The recommended concentration is 1.0  $\mu$ g/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

#### Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

## Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

# **Storage**

-20°C

#### **Precautions**

Anti-Human GRO- $\beta$  Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



# Anti-Human GRO-β Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Human GRO-β Antibody - Images