

Anti-Murine IL-10 Antibody
Catalog # ABG10195**Specification****Anti-Murine IL-10 Antibody - Product Information**

| | |
|-------------|------------|
| Application | WB, E |
| Reactivity | Mouse |
| Host | Rat |
| Clonality | Monoclonal |

Anti-Murine IL-10 Antibody - Additional Information**Preparation**

Produced in Sprague Dawley rats using highly pure (>98%) recombinant murine IL-10 as the immunizing antigen. This IgG2_A antibody was purified from cell culture by Protein G affinity chromatography.

WesternBlot

To detect mIL-10 by Western Blot analysis this antibody can be used at a concentration of 1.0-2.0 µg/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIL-10 is 2.0-4.0 ng/lane, under reducing or non-reducing conditions.

Sandwich

In a sandwich ELISA (assuming 100µl/well), a concentration of 4.0-8.0 µg/ml of this antibody will detect at least 1600 pg/ml of recombinant murine IL-10 when used with BioGems®; biotinylated antigen affinity purified anti-murine IL-10 (61-010BT) as the detection antibody at a concentration of approximately 0.5-1.0 µg/ml.

Neutralization

To yield one-half maximal inhibition **[ND₅₀]** of the biological activity of Murine IL-10 (0.5 ng/ml), a concentration of 3.0-5.0 µg/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Murine IL-10 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Murine IL-10 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Murine IL-10 Antibody - Images