

Anti-Murine IL-1 β Antibody
Catalog # ABG10243**Specification**

Anti-Murine IL-1 β Antibody - Product Information

Application	WB, IHC, E
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal

Anti-Murine IL-1 β Antibody - Additional Information**Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant mIL-1 β . Anti-Murine IL-1 β specific antibody was purified by affinity chromatography employing immobilized mIL-1 β matrix.

WesternBlot

To detect mIL-1 β by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 μ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIL-1 β is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect mIL-1 β by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems's Biotinylated Anti-Murine IL-1 β (61-001BBT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant mIL-1 β .

Immunohistochemistry

This antibody stained colchicine injected mouse brain (polymorph layer of the dentate gyrus) tissue. The primary antibody was incubated at 0.5 μ g/ml overnight at 4°C. This was followed by a peroxidase conjugated secondary antibody and then a fluorescein Tyramide Signal Amplification (TSA™) reagent. Optimal concentrations and conditions may vary. **Information and photo are courtesy of the Tissue Profiling group, SciLifeLab Stockholm.**

Neutralization

To yield one-half maximal inhibition [**ND**₅₀] of the biological activity of mIL-1 β (10 pg/ml), a concentration of 0.57-0.86 μ g/ml of this antibody is required.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Murine IL-1 β Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Murine IL-1 β Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Murine IL-1 β Antibody - Images