

Biotinylated Anti-Murine IL-5 Antibody

Catalog # ABG10289

Specification

Biotinylated Anti-Murine IL-5 Antibody - Product Information

Application WB, E
Reactivity Mouse
Host Rabbit
Clonality Polyclonal

Biotinylated Anti-Murine IL-5 Antibody - Additional Information

Preparation

Produced from sera of rabbits immunized with highly pure recombinant Murine IL-5. Anti-Murine IL-5 specific antibody was purified by affinity chromatography and then biotinylated.

WesternBlot

To detect Murine IL-5 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 mg/ml. When used in conjunction with compatible development reagents, the detection limit for recombinant Murine IL-5 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Murine IL-5 by sandwich ELISA (using 100 ml/well) a concentration of $0.25 - 1.0 \,\mu\text{g/ml}$ of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems' Polyclonal Anti- Murine IL-5 (61-005P) as a capture antibody, allows the detection of at least $0.2 - 0.4 \,\text{ng/well}$ of recombinant Murine IL-5.

Direct

To detect Murine IL-5 by direct ELISA (using 100 ml/well) a concentration of approximately 1.0 μ g/ml of this antibody is required. This biotinylated polyclonal antibody allows the detection of at least 0.2 – 0.4 ng/well of recombinant Murine IL-5.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Biotinylated Anti-Murine IL-5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.





Tel: 858.875.1900 Fax: 858.875.1999

Biotinylated Anti-Murine IL-5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Biotinylated Anti-Murine IL-5 Antibody - Images