

Anti-Rat IL-7 Antibody

Catalog # ABG10303

Specification

Anti-Rat IL-7 Antibody - Product Information

Application	WB, E
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal

Anti-Rat IL-7 Antibody - Additional Information

Preparation

Produced from sera of rabbits immunized with highly pure recombinant Rat IL-7. Anti-Rat IL-7 specific antibody was purified by affinity chromatography employing an immobilized Rat IL-7 matrix.

WesternBlot

To detect Rat IL-7 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 μ g/ml. When used in conjunction with compatible secondary reagents the detection limit for recombinant Rat IL-7 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Rat IL-7 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Rat IL-7 () as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Rat IL-7.

Neutralization

To yield one-half maximal inhibition [ND₅₀] of the biological activity of Rat IL-7 (5.0 ng/ml), a concentration of 0.14 - 0.21 μ g/ml of this antibody is required.

Storage -20°C

Precautions

Anti-Rat IL-7 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Rat IL-7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence



- Immunoprecipitation
- <u>Flow Cytomety</u>
 <u>Cell Culture</u>

Anti-Rat IL-7 Antibody - Images