

Anti-Human Leptin Antibody

Catalog # ABG10345

Specification

Anti-Human Leptin Antibody - Product Information

Application WB, E
Reactivity Human
Host Mouse
Clonality Monoclonal

Anti-Human Leptin Antibody - Additional Information

Preparation

Produced in BALB/c mice using highly pure (>98%) recombinant human Leptin as the immunizing antigen. This IgG1_K antibody was purified from cell culture by Protein A affinity chromatography.

WesternBlot

To detect hLeptin by Western Blot analysis this antibody can be used at a concentration of 0.20- $0.40 \mu g/ml$. Used in conjunction with compatible secondary reagents the detection limit for recombinant hLeptin is 2.0- $4.0 \mu g/lane$, under non-reducing conditions.

Sandwich

In a sandwich ELISA (assuming $100\mu\text{I/weII}$), a concentration of 2.0-4.0 µg/ml of this antibody will detect recombinant human Leptin when used with BioGems' biotinylated antigen affinity purified anti-human Leptin (60-204BT) as the detection antibody at a concentration of approximately 0.5-1.0 µg/ml.

Formulation

A sterile filtered antibody solution was lyophilized from PBS.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human Leptin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Human Leptin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot





• <u>Immunohistochemistry</u>

- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Human Leptin Antibody - Images