

Anti-Human Maspin Antibody

Catalog # ABG10350

Specification

Anti-Human Maspin Antibody - Product Information

Application WB, IHC, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

Anti-Human Maspin Antibody - Additional Information

Preparation

Produced from sera of rabbits immunized with highly pure recombinant Human Maspin. Anti-Human Maspin specific antibody was purified by affinity chromatography employing an immobilized Human Maspin matrix.

WesternBlot

To detect hMaspin by Western Blot analysis this antibody can be used at a concentration of $0.1 - 0.2 \,\mu\text{g/ml}$. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMaspin is $1.5 - 3.0 \,\text{ng/lane}$, under either reducing or non-reducing conditions.

Sandwich

To detect hMaspin by sandwich ELISA (using 100μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human Maspin (60-210BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMaspin.

Immunohistochemistry

This antibody stained formalin-fixed paraffin-embedded sections of human normal breast and human breast lobular carcinoma in situ. The recommended concentration is 7.8 ng/mL- 15.625 ng/mL with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Optimal results for these conditions were achieved without antigen retrieval. Optimal concentrations and conditions may vary.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human Maspin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.





Anti-Human Maspin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Human Maspin Antibody - Images