

## **Anti-Human MIA-2 Antibody**

Catalog # ABG10389

# Specification

# **Anti-Human MIA-2 Antibody - Product Information**

Application WB, IHC, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

### Anti-Human MIA-2 Antibody - Additional Information

#### **Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hMIA-2 (human MIA-2). Anti-Human MIA-2 specific antibody was purified by affinity chromatography employing immobilized hMIA-2 matrix.

#### WesternBlot

To detect hMIA-2 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2  $\mu$ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hMIA-2 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

#### Sandwich

To detect hMIA-2 by sandwich ELISA (using  $100\mu$ I/well antibody solution) a concentration of 0.5 - 2.0  $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human MIA-2 (60-220BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hMIA-2.

### **Immunohistochemistry**

This antibody stained formalin-fixed paraffin-embedded sections of human pancreas infiltrating ductal adenocarcinoma tissue. The recommended concentration is  $5.0~\mu g/ml$  with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with an alcohol-soluble AEC chromogen. Optimal results for these conditions were achieved without antigen retrieval. Optimal concentrations and conditions may vary.

#### **Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

# Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

#### **Storage**

-20°C

#### Precautions

Anti-Human MIA-2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



# **Anti-Human MIA-2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

**Anti-Human MIA-2 Antibody - Images**