

Anti-Murine MIP-17 Antibody

Catalog # ABG10409

Specification

Anti-Murine MIP-1y Antibody - Product Information

Application WB, E
Reactivity Mouse
Host Rabbit
Clonality Polyclonal

Anti-Murine MIP-17 Antibody - Additional Information

Preparation

Produced from sera of rabbits pre-immunized with highly pure recombinant Murine MIP- 1γ . Anti-Murine MIP- 1γ specific antibody was purified by affinity chromatography employing immobilized Murine MIP- 1γ matrix.

WesternBlot

To detect Murine MIP-1 γ by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2 μ g/ml. When used in conjunction with compatible secondary reagents, the detection limit for recombinant Murine MIP-1 γ is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect Murine MIP-1 γ by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.5 - 2.0 μ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Murine MIP-1 γ (61-101BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant Murine MIP-1 γ .

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Murine MIP- 1γ Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-Murine MIP-17 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides





- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-Murine MIP-1γ Antibody - Images