

Biotinylated Anti-Human NT-4 Antibody

Catalog # ABG10447

Specification

Biotinylated Anti-Human NT-4 Antibody - Product Information

Application Reactivity Host Clonality WB, E Human Goat Polyclonal

Biotinylated Anti-Human NT-4 Antibody - Additional Information

Preparation

Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hNT-4. Anti-Human NT-4 specific antibody was purified by affinity chromatography and then biotinylated.

WesternBlot

To detect hNT-4 by Western Blot analysis this antibody can be used at a concentration of $0.1 - 0.2 \mu$ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hNT-4 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Sandwich

To detect hNT-4 by sandwich ELISA (using 100 μ l/well antibody solution) a concentration of 0.25 – 1.0 μ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with BioGems' Polyclonal Anti-Human NT-4 (60-253P) as a capture antibody, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hNT-4.

Direct

To detect hNT-4 by direct ELISA (using 100 μ l/well antibody solution) a concentration of 0.25 – 1.0 μ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 – 0.4 ng/well of recombinant hNT-4.

Formulation

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile PBS containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml.

Storage -20°C

Precautions

Biotinylated Anti-Human NT-4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Biotinylated Anti-Human NT-4 Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Biotinylated Anti-Human NT-4 Antibody - Images