

### **Anti-Human PIGF-1 Antibody**

Catalog # ABG10463

### **Specification**

### **Anti-Human PIGF-1 Antibody - Product Information**

Application WB, IHC, E
Reactivity Human
Host Rabbit
Clonality Polyclonal

#### **Anti-Human PIGF-1 Antibody - Additional Information**

#### **Preparation**

Produced from sera of rabbits pre-immunized with highly pure (>98%) recombinant hPIGF. Anti-Human PIGF specific antibody was purified by affinity chromatography employing immobilized hPIGF matrix.

#### WesternBlot

To detect hPIGF by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2  $\mu g/ml$ . Used in conjunction with compatible secondary reagents the detection limit for recombinant hPIGF is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

#### Sandwich

To detect hPIGF by sandwich ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.5 - 2.0  $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human PIGF (60-274BT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hPIGF.

#### **Immunohistochemistry**

This antibody stained formalin-fixed, paraffin-embedded sections of human breast invasive ductal carcinoma. The recommended concentration is 2.0  $\mu$ g/ml with an overnight incubation at 4°C. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

#### **Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

## Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

#### **Storage**

-20°C

#### Precautions

Anti-Human PIGF-1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



# **Anti-Human PIGF-1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

**Anti-Human PIGF-1 Antibody - Images**