

Anti-Human Resistin Antibody

Catalog # ABG10484

Specification

Anti-Human Resistin Antibody - Product Information

Application WB, IHC, E
Reactivity Human
Host Mouse
Clonality Monoclonal

Anti-Human Resistin Antibody - Additional Information

Preparation

Produced in BALB/c mice using highly pure (>98%) recombinant human Resistin as the immunizing antigen. This $IgG < sub > 2b\kappa < / sub >$ antibody was purified from cell culture by Protein A affinity chromatography.

WesternBlot

To detect hResistin by Western Blot analysis this antibody can be used at a concentration of 0.20- $0.40 \mu g/ml$. Used in conjunction with compatible secondary reagents the detection limit for recombinant hResistin is 0.20- $0.40 \mu g/lane$, under non-reducing conditions.

Sandwich

In a sandwich ELISA (assuming 100μ I/well), a concentration of 4.0- $8.0~\mu$ g/ml of this antibody will detect at least 200 pg/ml of recombinant human Resistin when used with BioGems' biotinylated antigen affinity purified anti-human Resistin (60-289BT) as the detection antibody at a concentration of at least 0.5- $1.0~\mu$ g/ml.

Immunohistochemistry

This antibody stained formalin-fixed, paraffin-embedded sections of human normal placenta. The recommended concentration is $1.0~\mu g/ml$ with a two hour incubation at room temperature. An HRP-labeled polymer detection system was used with a DAB chromogen. Heat induced antigen retrieval with a pH 6.0 sodium citrate buffer is recommended. Optimal concentrations and conditions may vary.

Formulation

A sterile filtered antibody solution was lyophilized from PBS.

Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Storage

-20°C

Precautions

Anti-Human Resistin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



Anti-Human Resistin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-Human Resistin Antibody - Images