

# **Anti-Human SCF Antibody**

Catalog # ABG10489

## **Specification**

# **Anti-Human SCF Antibody - Product Information**

Application WB, IHC, E
Reactivity Human
Host Goat
Clonality Polyclonal

### **Anti-Human SCF Antibody - Additional Information**

#### **Preparation**

Produced from sera of goats pre-immunized with highly pure (>98%) recombinant hSCF. Anti-Human SCF specific antibody was purified by affinity chromatography employing immobilized hSCF matrix.

#### WesternBlot

To detect hSCF by Western Blot analysis this antibody can be used at a concentration of 0.1-0.2  $\mu$ g/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant hSCF is 1.5-3.0 ng/lane, under either reducing or non-reducing conditions.

#### Sandwich

To detect hSCF by sandwich ELISA (using 100  $\mu$ l/well antibody solution) a concentration of 0.5 - 2.0  $\mu$ g/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with BioGems' Biotinylated Anti-Human SCF (60-291GBT) as a detection antibody, allows the detection of at least 0.2 - 0.4 ng/well of recombinant hSCF.

#### **Immunohistochemistry**

This antibody stained CACO-2 cells. The primary antibody was incubated at 2.0  $\mu$ g/ml overnight at 4°C followed by a fluorescent labeled secondary antibody. Optimal concentrations and conditions may vary. Information and photo are courtesy of the Cell Profiling group, SciLifeLab Stockholm.

### **Neutralization**

To yield one-half maximal inhibition [<strong>ND</strong><span style="font-size: 16px;"><sub>50</sub></span>] of the biological activity of hSCF (10.00 ng/ml), a concentration of 0.008-0.012 µg/ml of this antibody is required.

# **Formulation**

A sterile filtered antibody solution was lyophilized from PBS, pH 7.2.

# Reconstitution

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

### **Storage**

-20°C

#### **Precautions**

Anti-Human SCF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.



Tel: 858.875.1900 Fax: 858.875.1999



# **Anti-Human SCF Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

**Anti-Human SCF Antibody - Images**