

Anti-DPYD antibody
Catalog # ABO10210**Specification**

Anti-DPYD antibody - Product Information

Application	WB, E
Primary Accession	A01749
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for DPYD detection. Tested with WB, Direct ELISA in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-DPYD antibody - Additional Information**Application Details**

Western blot, 0.1-0.5 µg/ml

 Direct ELISA, 0.1-0.5 µg/ml

Subcellular Localization

Cytoplasm.

Tissue Specificity

Found in most tissues with greatest activity found in liver and peripheral blood mononuclear cells.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E. coli-derived human DPYD recombinant protein (Position: A356-Y511).

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C; for one year. After r°Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and thawing.

Anti-DPYD antibody - Protein Information

Anti-DPYD antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-DPYD antibody - Images

Anti-DPYD antibody - Background

DPYD (Dihydropyrimidine Dehydrogenase), also called DPD, is an enzyme that in humans is encoded by the DPYD gene. The protein encoded by this gene is a pyrimidine catabolic enzyme and the initial and rate-limiting factor in the pathway of uracil and thymidine catabolism. The structure of the DPYD gene contains 23 exons spanning about 950 kb. Using somatic cell hybrid strategies, the DPYD gene is mapped to the centromeric region of chromosome 1 between 1p22 and 1q21. By fluorescence in situ hybridization, the DPYD gene is mapped to 1p22. The highest level of DPD was found in monocytes followed by that in lymphocytes, granulocytes, and platelets, whereas no significant activity of DPD could be detected in erythrocytes. The activity of DPD in peripheral blood mononuclear cells was intermediate between that observed in monocytes and lymphocytes. By cDNA microarray, Western blot analysis, and luciferase reporter assay, the transcription factor LSF was identified as a positive regulator of DPYD.