

Anti-CD7 Picoband Antibody
Catalog # ABO10221**Specification**

Anti-CD7 Picoband Antibody - Product Information

Application	IHC-P, FC, E
Primary Accession	P09564
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for CD7 detection. Tested with IHC-P, FCM, Direct ELISA in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CD7 Picoband Antibody - Additional Information

Gene ID 924

Other Names

T-cell antigen CD7, GP40, T-cell leukemia antigen, T-cell surface antigen Leu-9, TP41, CD7, CD7

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml
Flow Cytometry, 1-3^{1/4}g/1x10⁶ cells
Direct ELISA, 0.1-0.5 µg/ml

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E. coli-derived human CD7 recombinant protein (Position: A26-D172).

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C; for one year. After r°Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and thawing.

Anti-CD7 Picoband Antibody - Protein Information

Name CD7

Function

Not yet known.

Cellular Location

Membrane; Single-pass type I membrane protein.

Anti-CD7 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD7 Picoband Antibody - Images**Anti-CD7 Picoband Antibody - Background**

CD7 (Cluster of Differentiation 7) is a protein that in humans is encoded by the CD7 gene. This gene encodes a transmembrane protein which is a member of the immunoglobulin superfamily. This protein is found on thymocytes and mature T cells. It plays an essential role in T-cell interactions and also in T-cell/B-cell interaction during early lymphoid development.