

## **Anti-CD7 Picoband Antibody**

Catalog # ABO10221

## **Specification**

# **Anti-CD7 Picoband Antibody - Product Information**

Application IHC-P, FC, E
Primary Accession P09564
Host Reactivity Human
Clonality Polyclonal
Format Lyophilized

**Description** 

Rabbit IgG polyclonal antibody for CD7 detection. Tested with IHC-P, FCM, Direct ELISA in Human.

## Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

# **Anti-CD7 Picoband Antibody - Additional Information**

#### Gene ID 924

### **Other Names**

T-cell antigen CD7, GP40, T-cell leukemia antigen, T-cell surface antigen Leu-9, TP41, CD7, CD7

# **Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1  $\mu$ g/ml<br/>br><br/>Flow Cytometry, 1-3 $\hat{1}^{1}$ 4g/1x10<sup>6</sup> cells<br/>br><br/>br> Direct ELISA, 0.1-0.5  $\mu$ g/ml<br/>br>

#### **Contents**

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

## **Immuno**aen

E. coli-derived human CD7 recombinant protein (Position: A26-D172).

## **Cross Reactivity**

No cross reactivity with other proteins.

Storage At -20°C; for one year. After r°Constitution,

at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and

thawing.

# **Anti-CD7 Picoband Antibody - Protein Information**

Name CD7



#### **Function**

Transmembrane glycoprotein expressed by T-cells and natural killer (NK) cells and their precursors (PubMed:<a href="http://www.uniprot.org/citations/7506726" target="\_blank">7506726</a>). Plays a costimulatory role in T-cell activation upon binding to its ligand K12/SECTM1 (PubMed:<a href="http://www.uniprot.org/citations/10652336" target="\_blank">10652336</a>). In turn, mediates the production of cytokines such as IL-2 (PubMed:<a href="http://www.uniprot.org/citations/1709867" target="\_blank">1709867</a>). On resting NK-cells, CD7 activation results in a significant induction of interferon-gamma levels (PubMed:<a href="http://www.uniprot.org/citations/7506726" target="\_blank">7506726</a>).

#### **Cellular Location**

Membrane; Single-pass type I membrane protein.

#### **Tissue Location**

Expressed on T-cells and natural killer (NK) cells and their precursors.

## **Anti-CD7 Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **Anti-CD7 Picoband Antibody - Images**

# **Anti-CD7 Picoband Antibody - Background**

CD7 (Cluster of Differentiation 7) is a protein that in humans is encoded by the CD7 gene. This gene encodes a transmembrane protein which is a member of the immunoglobulin superfamily. This protein is found on thymocytes and mature T cells. It plays an essential role in T-cell interactions and also in T-cell/B-cell interaction during early lymphoid development.