

Anti-IL17E Picoband Antibody
Catalog # ABO10305**Specification**

Anti-IL17E Picoband Antibody - Product Information

Application	WB, E
Primary Accession	Q9H293
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for IL17E detection. Tested with WB, ELISA(Cap) in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-IL17E Picoband Antibody - Additional Information

Gene ID 64806

Other Names

Interleukin-25, IL-25, Interleukin-17E, IL-17E, IL25, IL17E

Application Details

Western blot, 0.1-0.5 µg/ml
 ELISA(Cap), 0.1-0.5 µg/ml

Subcellular Localization

Secreted.

Tissue Specificity

Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and trachea.

Contents

Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E. coli-derived human IL17E recombinant protein (Position: Y33-G177).

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C; for one year. After r° Constitution, at 4°C; for one month. It°Can also be aliquotted and stored frozen at -20°C; for a longer time. Avoid repeated freezing and

thawing.

Anti-IL17E Picoband Antibody - Protein Information

Name IL25

Synonyms IL17E

Function

Cytokine produced by various cells such as eosinophils, T- helper type 2 (Th2) cells or epithelial cells that plays a role in internal safety of adaptive immune responses by regulating cytokine production (PubMed:25821217, PubMed:15860795). Promotes and augments T- helper type 2 responses locally or systemically (PubMed:25821217). Exerts its activity via its receptor composed of IL17RA and IL17RB for signal transduction (By similarity). In turn, stimulates the JAK2- STAT5A pathway and promotes the secretion of type-2 associated cytokines including IL4, IL9 and IL13 (PubMed:25821217). Induces also the release of IL8, and IL6 from eosinophils through the combined activation of MAPK and NF-kappa-B pathways (PubMed:15860795). Inhibits the differentiation of T-helper (Th17) cells via the production of IL4, IL5 and IL13 (PubMed:11754819).

Cellular Location

Secreted.

Tissue Location

Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and trachea

Anti-IL17E Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IL17E Picoband Antibody - Images