

Anti-ADAM2 Picoband Antibody
Catalog # ABO10339

Specification

Anti-ADAM2 Picoband Antibody - Product Information

Application	WB
Primary Accession	Q99965
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Disintegrin and metalloproteinase domain-containing protein 2(ADAM2) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-ADAM2 Picoband Antibody - Additional Information

Gene ID 2515

Other Names

Disintegrin and metalloproteinase domain-containing protein 2, ADAM 2, Cancer/testis antigen 15, CT15, Fertilin subunit beta, PH-30, PH30, PH30-beta, ADAM2, FTNB

Calculated MW

82457 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Membrane; Single-pass type I membrane protein.

Tissue Specificity

Expressed specifically in spermatogenic cells in the seminiferous cells. Not detected in fetal tissues.

Protein Name

Disintegrin and metalloproteinase domain-containing protein 2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human ADAM2 (231-274aa WIDENKIATTGEANELLHTFLRWKTSYLVLRPHDVAFLLYREK), different from the related mouse and rat sequences by eleven amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-ADAM2 Picoband Antibody - Protein Information

Name ADAM2

Synonyms FTNB

Function

Sperm surface membrane protein that may be involved in sperm- egg plasma membrane adhesion and fusion during fertilization. Could have a direct role in sperm-zona binding or migration of sperm from the uterus into the oviduct. Interactions with egg membrane could be mediated via binding between its disintegrin-like domain to one or more integrins receptors on the egg. This is a non catalytic metalloprotease-like protein.

Cellular Location

Membrane; Single-pass type I membrane protein.

Tissue Location


Expressed specifically in spermatogenic cells in the seminiferous cells. Not detected in fetal tissues


Anti-ADAM2 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-ADAM2 Picoband Antibody - Images





Western blot analysis of ADAM2 expression in rat testis extract (lane 1), mouse testis extract (lane 2) and MCF-7 whole cell lysates (lane 3). ADAM2 at 100KD was detected using rabbit anti-ADAM2 Antigen Affinity purified polyclonal antibody (Catalog # ABO10339) at 0.5 μ g/mL. The blot was developed using chemiluminescence (ECL) method .

Anti-ADAM2 Picoband Antibody - Background

ADAM2 (A Disintegrin and Metalloproteinase Domain 2), also known as FTNB or PH30, is an enzyme that in humans is encoded by the ADAM2 gene. This gene encodes a member of the ADAM (a disintegrin and metalloprotease domain) family. Members of this family are membrane-anchored proteins structurally related to snake venom disintegrins, and have been implicated in a variety of biological processes involving cell-cell and cell-matrix interactions, including fertilization, muscle development, and neurogenesis. This member is a subunit of an integral sperm membrane glycoprotein called fertilin, which plays an important role in sperm-egg interactions. This gene is mapped to 8p11.2.