

Anti-Annexin VIII Picoband Antibody
Catalog # ABO10341**Specification**

Anti-Annexin VIII Picoband Antibody - Product Information

Application	WB
Primary Accession	P13928
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Annexin A8(ANXA8) detection. Tested with WB in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Annexin VIII Picoband Antibody - Additional Information

Gene ID 653145;728113

Other Names

Annexin A8 {ECO:0000312|HGNC:HGNC:546}, Annexin VIII, Annexin-8, Vascular anticoagulant-beta, VAC-beta, ANXA8 (HGNC:546), ANX8

Calculated MW

36881 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human

Protein Name

Annexin A8

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human Annexin VIII (20-61aa HFNPDPDAETLYKAMKGIGTNEQAIIDVLTKRSTNRQQLAK), different from the related mouse and rat sequences by one amino acid.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-Annexin VIII Picoband Antibody - Protein Information

Name ANXA8 ([HGNC:546](#))

Synonyms ANX8

Function

This protein is an anticoagulant protein that acts as an indirect inhibitor of the thromboplastin-specific complex, which is involved in the blood coagulation cascade.

Anti-Annexin VIII Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Annexin VIII Picoband Antibody - Images




Figure 1. Western blot analysis of Annexin VIII using anti-Annexin VIII antibody (ABO10341). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. lane 1: HELA whole cell lysates, lane 2: A549 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90

minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Annexin VIII antigen affinity purified polyclonal antibody (Catalog # ABO10341) at 0.5 μ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Annexin VIII at approximately 37KD. The expected band size for Annexin VIII is at 37KD.

Anti-Annexin VIII Picoband Antibody - Background

ANXA8 is also known as ANNEXIN VIII. This gene encodes a member of the annexin family of evolutionarily conserved Ca²⁺ and phospholipid binding proteins. The encoded protein may function as an anticoagulant that indirectly inhibits the thromboplastin-specific complex. Overexpression of this gene has been associated with acute myelocytic leukemia. A highly similar duplicated copy of this gene is found in close proximity on the long arm of chromosome 10.