

**Anti-NDRG3 Picoband Antibody**  
**Catalog # ABO10345****Specification**

---

**Anti-NDRG3 Picoband Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	<a href="#">Q9UGV2</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Protein NDRG3(NDRG3) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-NDRG3 Picoband Antibody - Additional Information**

**Gene ID** 57446

**Other Names**

Protein NDRG3, N-myc downstream-regulated gene 3 protein, NDRG3

**Calculated MW**

41409 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat  
Western blot, 0.1-0.5 µg/ml, Human

**Tissue Specificity**

Ubiquitous. Highly expressed in brain. .

**Protein Name**

Protein NDRG3

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

**Immunogen**

E.coli-derived human NDRG3 recombinant protein (Position: L181-R239). Human NDRG3 shares 93.1% and 94.9% amino acid (aa) sequence identity with mouse and rat NDRG3, respectively.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

Storage

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

### Anti-NDRG3 Picoband Antibody - Protein Information

**Name** NDRG3

**Tissue Location**

Ubiquitous. Highly expressed in brain.

### Anti-NDRG3 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-NDRG3 Picoband Antibody - Images




Figure 1. Western blot analysis of NDRG3 using anti- NDRG3 antibody (ABO10345). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: COLO320 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti- NDRG3 antigen

affinity purified polyclonal antibody (Catalog # ABO10345) at 0.5  $\mu$ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for NDRG3 at approximately 41KD. The expected band size for NDRG3 is at 41KD.




Figure 2. IHC analysis of NDRG3 using anti- NDRG3 antibody (ABO10345).NDRG3 was detected in paraffin-embedded section of mouse brain tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti- NDRG3 Antibody (ABO10345) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




Figure 3. IHC analysis of NDRG3 using anti- NDRG3 antibody (ABO10345).NDRG3 was detected in paraffin-embedded section of rat brain tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1  $\mu$ g/ml rabbit anti- NDRG3 Antibody (ABO10345) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




Figure 4. IHC analysis of NDRG3 using anti- NDRG3 antibody (ABO10345).NDRG3 was detected in paraffin-embedded section of human intestinal cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu$ g/ml rabbit anti- NDRG3 Antibody (ABO10345) overnight at 4 $^{\circ}$ C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 $^{\circ}$ C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




Figure 5. IHC analysis of NDRG3 using anti- NDRG3 antibody (ABO10345).NDRG3 was detected in paraffin-embedded section of human mammary cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu$ g/ml rabbit anti- NDRG3 Antibody (ABO10345) overnight at 4 $^{\circ}$ C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 $^{\circ}$ C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

#### **Anti-NDRG3 Picoband Antibody - Background**

In studies of the NDRG gene family in the human, novel mouse sequences of Ndr2 and Ndr3 were used to search the human genome databases. This confirmed the existence of a human NDRG3 gene, homologous to mouse Ndr3. The 3 known human NDRG proteins show considerable homology, with 54% amino acid sequence identity between NDRG1 and NDRG2, 67% between NDRG1 and NDRG3, and 58% between NDRG2 and NDRG3. The NDRG3 gene was represented by 86 ESTs and a genomic clone from chromosome 20q11.21-q11.23. This provisional chromosomal localization was confirmed by electronic PCR.