

Anti-GJB2 Antibody
Catalog # ABO10518**Specification**

Anti-GJB2 Antibody - Product Information

Application	WB
Primary Accession	P29033
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Gap junction beta-2 protein(GJB2) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-GJB2 Antibody - Additional Information

Gene ID 2706

Other Names

Gap junction beta-2 protein, Connexin-26, Cx26, GJB2

Calculated MW

26215 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse

Subcellular Localization

Cell membrane; Multi-pass membrane protein. Cell junction, gap junction.

Protein Name

Gap junction beta-2 protein

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human GJB2(1-16aa, MDWGTLQTILGGVNH), different from the related mouse and rat sequences by two amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the connexin family. Beta-type (group I) subfamily.

Anti-GJB2 Antibody - Protein Information**Name** GJB2**Function**

Structural component of gap junctions (PubMed:16849369, PubMed:17551008, PubMed:19340074, PubMed:19384972, PubMed:21094651, PubMed:26753910). Gap junctions are dodecameric channels that connect the cytoplasm of adjoining cells. They are formed by the docking of two hexameric hemichannels, one from each cell membrane (PubMed:17551008, PubMed:19340074, PubMed:21094651, PubMed:26753910). Small molecules and ions diffuse from one cell to a neighboring cell via the central pore (PubMed:16849369, PubMed:19384972, PubMed:21094651).


Cellular Location


Cell membrane; Multi-pass membrane protein. Cell junction, gap junction. Note=Colocalizes with GJB4 at gap junction plaques in the cochlea. {ECO:0000250|UniProtKB:Q00977}

Anti-GJB2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-GJB2 Antibody - Images



Anti-GJB2 Antibody - Background

Connexin26(CX26), also known as GAP junction protein, beta2, GJB2. Gap junctions were first characterized by electron microscopy as regionally specialized structures on plasma membranes of contacting adherent cells. These structures were shown to consist of cell-to-cell channels. Proteins, called connexins, purified from fractions of enriched gap junctions from different tissues differ. The 3-prime untranslated region of the CX26 transcript contains a putative mRNA instability sequence. The deduced 226-amino acid protein has a calculated molecular mass of about 26 kD. CX26 shares 92.5% identity with rat Cx26. connexin 26(GJB2) is assigned to human chromosome 13q11-q12 .Connexin 26 regulates epidermal barrier and wound remodeling and promotes psoriasiform response. Connexin 26 gene(GJB2) mutation modulates the severity of hearing loss associated with the 1555A-G mitochondrial mutation.