10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-Cardiac FABP Antibody

**Catalog # ABO10662**

## Specification

### Anti-Cardiac FABP Antibody - Product Information

| | |
|---|---|
| Application | **WB, IHC-P, IHC-F** |
| Primary Accession | P05413 |
| Host | **Rabbit** |
| Reactivity | **Human, Mouse, Rat** |
| Clonality | **Polyclonal** |
| Format | **Lyophilized** |

**Description**
Rabbit IgG polyclonal antibody for Fatty acid-binding protein, heart(FABP3) detection. Tested with WB, IHC-P, IHC-F in Human;Mouse;Rat.

**Reconstitution**
Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

### Anti-Cardiac FABP Antibody - Additional Information

**Gene ID** 2170

**Other Names**
Fatty acid-binding protein, heart, Fatty acid-binding protein 3, Heart-type fatty acid-binding protein, H-FABP, Mammary-derived growth inhibitor, MDGI, Muscle fatty acid-binding protein, M-FABP, FABP3, FABP11, MDGI

**Calculated MW**
14858 MW KDa

**Application Details**
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat<br> <br>Immunohistochemistry(Frozen Section), 0.5-1 µg/ml, Human, -<br>Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat<br>

**Subcellular Localization**
Cytoplasm.

**Protein Name**
Fatty acid-binding protein, heart

**Contents**
Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

**Immunogen**
A synthetic peptide corresponding to a sequence at the C-terminus of human FABP3 (119-133aa THGTAVCTRTYEKEA), different from the related mouse and rat sequences by three amino acids.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**
No cross reactivity with other proteins

| | |
|---|---|
| Storage | **At -20˚C for one year. After r˚Constitution, at 4˚C for one month. It˚Can also be aliquotted and stored frozen at -20˚C for a longer time.Avoid repeated freezing and thawing.** |

## Anti-Cardiac FABP Antibody - Protein Information

**Name** FABP3

**Synonyms** FABP11, MDGI

**Function**
FABPs are thought to play a role in the intracellular transport of long-chain fatty acids and their acyl-CoA esters.

**Cellular Location**
Cytoplasm.

## Anti-Cardiac FABP Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## Anti-Cardiac FABP Antibody - Images

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999




Western blot analysis of Cardiac FABP expression in rat liver extract (lane 1), mouse cardiac muscle extract (lane 2), HELA whole cell lysates (lane 3) and MCF-7 whole cell lysates (lane 4). Cardiac FABP at 15KD was detected using rabbit anti- Cardiac FABP Antigen Affinity purified polyclonal antibody (Catalog # ABO10662) at 0.5 Î¼g/mL. The blot was developed using chemiluminescence (ECL) method .



Cardiac FABP was detected in paraffin-embedded sections of rat cardiac muscle tissues using rabbit anti- Cardiac FABP Antigen Affinity purified polyclonal antibody (Catalog # ABO10662) at 1 Î¼g/mL. The immunohistochemical section was developed using SABC method .



Cardiac FABP was detected in paraffin-embedded sections of mouse skeletal muscletissues using rabbit anti- Cardiac FABP Antigen Affinity purified polyclonal antibody (Catalog # ABO10662) at 1 Î¼g/mL. The immunohistochemical section was developed using SABC method .

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Cardiac FABP was detected in paraffin-embedded sections of human intestinal cancer tissues using rabbit anti- Cardiac FABP Antigen Affinity purified polyclonal antibody (Catalog # ABO10662) at 1 ??g/mL. The immunohistochemical section was developed using SABC method .

**Anti-Cardiac FABP Antibody - Background**

Heart-type fatty acid binding protein(hFABP) also known as mammary-derived growth inhibitor is a protein that in humans is encoded by the FABP3 gene. The intracellular fatty acid-binding proteins(FABPs) belongs to a multigene family. Fatty acid-binding protein 3 gene contains four exons and its function is to arrest growth of mammary epithelial cells. This gene is also a candidate tumor suppressor gene for human breast cancer. Cardiac-type fatty acid-binding protein(cFABP) from human heart muscle of three individuals was isolated and characterized as pI 5.3-cFABP.