

Anti-IGF-1 Antibody

Catalog # ABO10699

Specification

Anti-IGF-1 Antibody - Product Information

ApplicationWB, IHC-PPrimary AccessionP05019HostRabbitReactivityHuman, Mouse, RatClonalityPolyclonalFormatLyophilizedDescriptionRabbit lgG polyclonal antibody for Insulin-like growth factor I(IGF1) detection. Tested with WB, IHC-P in Human; Mouse; Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-IGF-1 Antibody - Additional Information

Gene ID 3479

Other Names Insulin-like growth factor I, IGF-I, Mechano growth factor, MGF, Somatomedin-C, IGF1, IBP1

Calculated MW 21841 MW KDa

Application Details Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, Mouse, By Heat

 Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization Secreted.

Protein Name Insulin-like growth factor I(IGF-I)

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence in the middle region of human IGF-1(80-96 aa GSSSRRAPQTGIVDECC), different from the mouse and rat sequences by one amino acid.

Purification Immunogen affinity purified.

Cross Reactivity



No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Sequence Similarities Belongs to the insulin family.

Anti-IGF-1 Antibody - Protein Information

Name IGF1 (HGNC:5464)

Function

The insulin-like growth factors, isolated from plasma, are structurally and functionally related to insulin but have a much higher growth-promoting activity. May be a physiological regulator of [1-14C]- 2-deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblasts. Stimulates glucose transport in bone-derived osteoblastic (PyMS) cells and is effective at much lower concentrations than insulin, not only regarding glycogen and DNA synthesis but also with regard to enhancing glucose uptake. May play a role in synapse maturation (PubMed:21076856, PubMed:24132240). Ca(2+)-dependent exocytosis of IGF1 is required for sensory perception of smell in the olfactory bulb (By similarity). Acts as a ligand for IGF1R. Binds to the alpha subunit of IGF1R, leading to the activation of the intrinsic tyrosine kinase activity which autophosphorylates tyrosine residues in the beta subunit thus initiating a cascade of down-stream signaling events leading to activation of the PI3K-AKT/PKB and the Ras-MAPK pathways. Binds to integrins ITGAV:ITGB3 and ITGA6:ITGB4. Its binding to integrins and subsequent ternary complex formation with integrins and IGFR1 are essential for IGF1 signaling. Induces the phosphorylation and activation of IGFR1, MAPK3/ERK1, MAPK1/ERK2 and AKT1 (PubMed:19578119, PubMed:22351760, PubMed:23243309, PubMed:23696648). As part of the MAPK/ERK signaling pathway, acts as a negative regulator of apoptosis in cardiomyocytes via promotion of STUB1/CHIP-mediated ubiquitination and degradation of ICER-type isoforms of CREM (By similarity).

Cellular Location Secreted {ECO:0000250|UniProtKB:P05017}.

Anti-IGF-1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>



Anti-IGF-1 Antibody - Images



Anti-IGF-1 antibody, ABO10699, Western blottingWB: Recombinant Human IGF-1 Protein 2.5ng



Anti-IGF-1 antibody, ABO10699, IHC(P)IHC(P): Human Mammary Cancer Tissue

Anti-IGF-1 Antibody - Background

Insulin-like growth factor 1(IGF-1) also known as somatomedin C or mechano growth factor is a protein that in humans is encoded by the IGF1 gene. IGF-1 is a hormone similar in molecular structure to insulin. It plays an important role in childhood growth and continues to have anabolic effects in adults. A synthetic analog of IGF-1, mecasermin is used for the treatment of growth failure. IGF-1 consists of 70 amino acids in a single chain with three intramolecular disulfide bridges. IGF-1 has a molecular weight of 7649 daltons.Justice et al.(1990) placed the mouse IGF1 gene on chromosome 10.